





Two fold (Double) eqn of two straight lines:-



$$ax + by = 0$$

$$ax + by + c = 0$$

The two lines $L_1: a_1x + b_1y + c_1 = 0$

& $L_2: a_2x + b_2y + c_2 = 0$

are

a) $|| \Leftrightarrow \frac{a_1}{a_2} = \frac{b_1}{b_2}$

b) The same if $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$

$$\tan \theta = \frac{m_2 - m_1}{1 + m_1 m_2}$$


~~$y = m_1x + c$~~ θ $y = m_2x + c_2$

* The distance bet. $P_1(x_1, y_1)$ & $P_2(x_2, y_2)$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

* The distance from $P_1(x_1, y_1)$ to $L: ax + by + c = 0$ is

$$d = \frac{ax_1 + by_1 + c}{\sqrt{a^2 + b^2}}$$



EX: Find the line $\parallel L$ to $4x - 5y + 1 = 0$

passing thr $(1, 2)$

Soln

Any line $\parallel L$ to (i) is given by $4x - 5y + K = 0$
 $(1, 2)$

$$4 - 10 + K = 0$$

$$K = 6$$

$$4x - 5y + 6 = 0 \quad \#$$

Subject. _____

Date. _____

The homogeneous eqn. of 2nd degree in x, y represents two lines passing thr 0. $ax^2 + 2hxy + by^2 = 0 \rightarrow$ \star
 بالقسمة على x^2

$$b\left(\frac{y}{x}\right)^2 + 2h\left(\frac{y}{x}\right) + a = 0$$

$$\frac{y}{x} = \frac{-2h \pm \sqrt{4h^2 - 4ab}}{2b}$$

$$\frac{y}{x} = \frac{-h \pm \sqrt{h^2 - ab}}{b}$$

$$y = \left(\frac{-h \pm \sqrt{h^2 - ab}}{b} \right) x$$

$$y = m_1 x \quad \& \quad y = m_2 x$$

$$m_1 = \frac{-h + \sqrt{h^2 - ab}}{b}$$

$$m_2 = \frac{-h - \sqrt{h^2 - ab}}{b}$$

provided $h^2 - ab \geq 0$

في تمثل خطين لا زلز المميز يكون
 أكبر من أو يساوي صفر

EX: Find the two lines represented by

$$y^2 - 7xy + 10x^2 = 0$$

$$(y - 2x)(y - 5x) = 0$$

$$y = 2x \quad \& \quad y = 5x$$

Note:- $m_1 + m_2 = \frac{-2h}{b}$

$$m_1 m_2 = \frac{a}{b}$$

The angle θ bet. the two lines *

$$\tan \theta = \frac{m_2 - m_1}{1 + m_1 m_2} = \frac{\sqrt{(m_2 - m_1)^2}}{1 + m_1 m_2}$$

$$= \frac{\sqrt{(m_1 + m_2)^2 - 4m_1 m_2}}{1 + m_1 m_2}$$

$$\tan \theta = \frac{\sqrt{\frac{4h^2}{b^2} - 4\frac{a}{b}}}{1 + \frac{a}{b}} = \frac{2\sqrt{h^2 - ab}}{a + b}$$

$\therefore a = 10, b = 1, h = \frac{7}{2}$

$$\tan \theta = \frac{2\sqrt{\frac{49}{4} - 10}}{11} = \frac{3}{11}$$


Subject. _____

Date. _____

The double eqn of the bisectors of \angle of \odot

$$d_1 = \pm d_2 \quad \text{Circalibi}$$

$$\left(\frac{y - m_1 x}{\sqrt{1 + m_1^2}} \right) = \pm \frac{y - m_2 x}{\sqrt{1 + m_2^2}}$$



$$\left(\frac{y - m_1 x}{\sqrt{1 + m_1^2}} - \frac{y - m_2 x}{\sqrt{1 + m_2^2}} \right) \left(\frac{y - m_1 x}{\sqrt{1 + m_1^2}} + \frac{y - m_2 x}{\sqrt{1 + m_2^2}} \right) = 0$$

$$\frac{(y - m_1 x)^2}{1 + m_1^2} - \frac{(y - m_2 x)^2}{1 + m_2^2} = 0$$

$$(1 + m_2^2)(y^2 + m_1^2 x^2 - 2m_1 xy) - (1 + m_1^2)(y^2 - 2m_2 xy + m_2^2 x^2) = 0$$

$$(m_1 + m_2)(x^2 - y^2) + 2(m_1 m_2 - 1)xy = 0$$

$$-\frac{2h}{b}(x^2 - y^2) + 2\left(\frac{a}{b} - 1\right)xy = 0 \quad (*b$$

$$h(x^2 - y^2) = (a - b)xy$$

$$\therefore \frac{x^2 - y^2}{a - b} = \frac{xy}{h}$$


Subject: _____

XI-Prove that the two lines thro. and make an angle
 with the line $x-y=0$ is given by
 $x^2 \pm 2xy \sec 2\alpha + y^2 = 0$

$$ax^2 + 2hxy + by^2 = 0$$

$$x-y=0 \Rightarrow x+y=0$$

$$x^2 - y^2 = 0 \text{ ----- (1)}$$



But the eqn of the bisectors

$$x^2 - y^2 = \frac{a-b}{h} xy \text{ ----- (2)}$$

$$(1) = (2)$$

$$\frac{a-b}{h} = 0 \Rightarrow \boxed{a=b}$$

$$\tan 2\alpha = \frac{2\sqrt{h^2 - a^2}}{2a}$$

$$a^2 \tan^2 2\alpha = h^2 - a^2$$

$$h^2 = a^2 (\tan^2 2\alpha + 1)$$

$$\boxed{h = \pm a \sec 2\alpha}$$

$$ax^2 \pm 2a \sec 2\alpha \cdot xy + ay^2 = 0$$